

and maintained grass lawn on the north side. A query of VertNet (www.vertnet.org) yielded no unpublished records in Madison County. The nearest known record is ca. 13.6 km to the south-southeast in Chester County (Cornell University Museum of Vertebrates [CUMV] R-0002319).

LEE J. BARTON, Freed-Hardeman University, 151 E Main Street, Henderson, Tennessee 38340, USA; e-mail: lbarton@fhu.edu.

HALDEA STRIATULA (Rough Earthsnake). USA: TEXAS: KARNES Co.: Joe Gulley Park, Kenedy (28.81566°N, 97.86218°W; WGS 84). 5 March 2022. Drew R. Davis. Verified by Travis J. LaDuc. Biodiversity Collections, The University of Texas at Austin (TNHC 116650 [DRD 9026]). Subadult male (110 mm SVL, 19 mm tail length [incomplete], 1.1 g) collected under a rock at 1600 h. This specimen represents a new county record (Dixon 2013. *Amphibians and Reptiles of Texas: with Keys, Taxonomic Synopses, Bibliography, and Distribution Maps*. Third Edition Revised and Updated. Texas A&M University Press, College Station, Texas. viii + 447 pp.) and fills a gap along the southwestern edge of the species' recognized distribution in the state. *Haldea striatula* is known from five bordering counties (Dixon 2013, *op. cit.*), and the nearest known specimen record is ca. 56.0 km to the northwest in Wilson County (TNHC 100684). This specimen was collected under a Texas Parks and Wildlife Scientific Collecting Permit (SPR-1018-294) and an approved IACUC protocol (UTRGV AUP #18-28), both issued to DRD.

BENJAMIN W. GENTER, Columbus North High School, 1400 25th Street, Columbus, Indiana 47201, USA (e-mail: benwgenter@gmail.com);

DREW R. DAVIS, School of Earth, Environmental, and Marine Sciences, The University of Texas Rio Grande Valley, 1 W University Boulevard, Brownsville, Texas 78520, USA and Biodiversity Collections, Department of Integrative Biology, The University of Texas at Austin, 10100 Burnet Road, PRC 176-R4000, Austin, Texas 78758, USA (e-mail: drew.davis@utrgv.edu).

LYCODON PAUCIFASCIATUS (Rendahl's Big-tooth Snake). VIETNAM: QUANG NGAI PROVINCE: BA TO DISTRICT: Cao Muon Mountain, Ba Vinh Commune (ca. 14.80192°N, 108.68172°E; WGS 84), 400 m elev. 6 May 2020. P. M. Nguyen. Verified by Indraneil Das. Herpetological Collection, Duy Tan University (DTU 439; photo voucher). Individual found at 2000 h foraging in stream; surrounding habitat was secondary forest.

KIEN GIANG PROVINCE: PHU QUOC CITY: Bai Thom Commune within Phu Quoc National Park (ca. 10.36616°N, 103.99765°E; WGS 84), 150 m elev. 15 June 2021. T.T. Vo. Verified by Indraneil Das. DTU 462 (photo voucher). Individual found at 1940 h in under rock near stream; surrounding habitat was secondary forest.

DAK LAK PROVINCE: EA KRA DISTRICT: Ea So Commune within Ea So Nature Reserve (ca. 12.94740°N, 108.47697°E; WGS 84), 570 m. elev. 20 June 2021. L. T. Nguyen. Verified by Indraneil Das. DTU 463 (photo voucher). Individual found at 2120 h crawling on branch 2 m above ground; surrounding habitat was secondary forest mixed with shrubs.

First records for Quang Ngai, Dak Lak and Kien Giang provinces, as well as third record for Vietnam. The species is endemic to central Vietnam; the locality in Quang Ngai is ca. 174 airline km northwest of the nearest population in Thua Luu Distric, Thua Thien-Hue Province, Vietnam (Vogel et al. 2009. *Trop Zool.* 22:131–182). We are grateful to P. M. Nguyen (Quang Ngai), T.T. Vo (Kien Giang), and L. T. Nguyen (Dak Lak) for providing photos and information.

PHUONG KY NGOC NGUYEN, Da Lat University, Da Lat, Lam Dong (e-mail: nguyenphuongdlu0310@gmail.com); **TANG VAN DUONG**, Vietnam National Museum of Nature, Vietnam Academy of Science and Technology, Hanoi, Vietnam (e-mail: tangdv@vnmn.vast.vn); **TAN VAN NGUYEN**, Save Vietnam's Wildlife, Nho Quan, Ninh Binh, Vietnam (e-mail: tanvannguyen150991@gmail.com).

NERODIA FLORIDANA (Florida Green Watersnake). USA: FLORIDA: SANTA ROSA Co.: Escribano Point Wildlife Management Area (30.50926°N, 86.99492°W; WGS 84). 8 March 2022. Kameron Burgess, Samantha Dillon, and Wynn Hoskins. Verified by Coleman M. Sheehy III. Florida Museum of Natural History, University of Florida (UF 193001; photo voucher). Juvenile found on a flooded dirt trail in wet pine flatwoods. New county record (Krysko et al. 2019. *Amphibians and Reptiles of Florida*. University Press of Florida, Gainesville, Florida. 706 pp.) and extends its range 67 km west of the nearest record in Walton County, Florida (UF 53844). The record is only 20 km east-southeast of the nearest *Nerodia cyclopion* (Mississippi Green Watersnake) record from near the mouth of the Escambia River in Escambia County, Florida (Auburn University Museum of Natural History [AUM] 2132), but is separated from it by brackish Blackwater and Escambia bays and Garcon Point, a peninsula of land separating the bays.

KAMERON BURGESS and **SAMANTHA DILLON**, The Longleaf Alliance, 8831 Whiting Field Circle, Milton, Florida 32570, USA; **KEVIN M. ENGE**, Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission, 1105 S.W. Willison Road, Gainesville, Florida 32601, USA (e-mail: kevin.enge@myfwc.com).

REGINA GRAHAMII (Graham's Crayfish Snake). USA: KANSAS: ELK Co.: Tributary of Grouse Creek on Youngmeyer Ranch, 0.74 km N Wiededman Rd (37.32030°N, 96.30580°W; NAD 83). 9 June 2020. Krista J. Ward, Thomas M. Luhring, Jake T. Wright, and Christine S. Streid. Verified by Dexter R. Mardis. University of Kansas Digital Archives (KUDA 13749; photo voucher). A single individual (31 cm SVL) found in a shallow (0.12 m depth) ephemeral stream pool in an upland pasture. New county record (Taggart 2020. *Kansas Herpetofaunal Atlas: An On-line Reference*. Electronic Database available at <https://webapps.fhsu.edu/ksherp>. Accessed 20 Jun 2020). This record fills in a gap in the distribution of this species in southeast Kansas. It has been recorded from the adjacent Butler, Cowley, Republic, Montgomery, Wilson, and Greenwood counties (Taggart 2020, *op. cit.*). The closest known record is 15.94 km to the west in Butler County (Biodiversity Institute, University of Kansas [KU] 318692).

KRISTA J. WARD (e-mail: kristajoy.ward@gmail.com), **CHRISTINE S. STREID** (e-mail: csstreid@shockers.wichita.edu), **JAKE T. WRIGHT** (e-mail: jtwright3@shockers.wichita.edu), and **THOMAS M. LUHRING**, Department of Biology, Wichita State University, 1845 Fairmount Street, Wichita, Kansas 67260, USA (e-mail: thomas.luhring@wichita.edu).

SONORA SEMIANNULATA (Western Groundsnake). USA: ARIZONA: GREENLEE Co.: 13.6 km south of Franklin (32.54656°N, 109.05886°W; WGS 84), 1244 m elev. 29 August 2021. Jaron. P. Lopez. Verified by Brian K. Sullivan. Arizona State University (ASU HP000614; photo voucher). The snake was photographed moving on the surface and not captured. Adult with 34 black bands and orange dorsum. New county record (Holycross and Mitchell [eds] 2020. *Snakes of Arizona*. ECO Wear & Publishing, Rodeo, New Mexico. 860 pp.) and fills a gap between records ca.