

Catalogue of American Amphibians and Reptiles.

CALDWELL, JANALEE P. 1982. *Hyla gratiosa**Hyla gratiosa* LeConte
Barking treefrog

Hyla gratiosa LeConte, 1856:146. Type-locality, "lower country of Georgia," restricted to "Liberty County, Georgia" by Schmidt (1953:71). Holotype, Acad. Natur. Sci. Philadelphia 2089 according to Malnate (1971:351; examined by author). Cochran's (1961:54) listing of six "cotypes" from Riceboro, Liberty County, in the National Museum of Natural History is puzzling, for LeConte mentioned only three specimens.

Epedaphus gratiosus: Cope, 1885:383.

• CONTENT. No subspecies are recognized.

• DEFINITION AND DIAGNOSIS. A member of the *Hyla cinerea* species group (Blair, 1959), *gratiosa* is the largest *Hyla* in the United States: females 50–70 mm snout–vent length, males 49–70 mm. Characteristics include: stout forelimbs, fingers webbed at base with large terminal discs; toes about one-half webbed; skin evenly granulated over entire body; thoracic fold present. The dorsal color is variable bright green to dark olive green with evenly spaced darker round spots encircled with black, sometimes indistinct. An irregular white stripe extends from the upper lip to the groin.

Hyla cinerea is similar to *H. gratiosa* in size and coloration but its body is more slender, its lateral white line is more sharply defined, and it never has dark, round spots on the dorsum. There is no overlap of the ratio of head width to tibio-fibula length in *H. gratiosa* and *H. cinerea* (Gerhardt et al., 1980). *Hyla andersoni* is smaller and has a distinct mask. *Hyla squirella* may be green with a light lateral stripe, but small size and generally brown or variable coloration distinguish it from *gratiosa*.

• DESCRIPTIONS. For adults, see Cope (1889) and Wright and Wright (1949); tadpoles, Wright (1929, 1932) and Altig (1970, 1972); eggs, Livezey and Wright (1947), Wright (1932), and Wright and Wright (1949). For keys to adults, see Carr and Goin (1955), Mount (1975), and Blair et al. (1957); tadpoles, Altig (1970) and Wright (1929); eggs, Livezey and Wright (1947).

• ILLUSTRATIONS. Black-and-white illustrations of adults are in LeConte (1856), Cope (1889), Wright (1932), Wright and Wright (1949), Carr and Goin (1955), Blair et al. (1957), and Mount (1975). Color illustrations are in Conant (1975) and Behler and King (1979). A hybrid *gratiosa* × *cinerea* is figured by Lee (1968a). For pre-

metamorphic stages, see Altig (1972), Livezey and Wright (1947), Wright (1932), and Wright and Wright (1949). Altig and Pace (1974) figure an electron photo-micrograph of a tadpole's labial tooth. Some skeletal and soft anatomy is illustrated in Cope (1889), Gaudin (1974), and Hilton (1954).

• DISTRIBUTION. *Hyla gratiosa* occurs on the Atlantic and Gulf coastal plains from the Florida parishes in Louisiana to southern Florida and north to Virginia. The species occurs in the Alabama piedmont, but is absent from most of the Georgia piedmont. Disjunct populations are reported from Kentucky (Monroe and Taylor, 1972; Monroe and Giannini, 1977), Tennessee (Jacob, 1980; Rossman, 1958), but a colony in New Jersey (Black and Gosner, 1958) may be extinct (K. L. Gosner, pers. comm.). Additional comments on distribution are by deRageot et al. (1969) and Mount (1975). *Hyla gratiosa* is absent from the majority of the Atlantic coast barrier islands (Gibbons and Coker, 1978; Martof, 1963) and from the Florida barrier islands as well (Blaney, 1971).

• FOSSIL RECORD. Chantell (1964) referred ilia from the Miocene Norden Bridge Local Fauna, Lower Valentine formation, Brown County, Nebraska, to *Hyla* cf. *H. gratiosa*.

• PERTINENT LITERATURE. Blair (1959) included *Hyla gratiosa*, *H. cinerea*, and *H. andersoni* in the *cinerea* species group based on similarity of morphology and calls. Hybridization of *gratiosa* in nature and in the laboratory is reported in numerous papers: Blair (1959), Fortman and Altig (1973, 1974), Gerhardt (1974), Gerhardt et al. (1980), Lee (1968a, 1968b), Mecham (1960, 1965), Moore (1955), and Pyburn and Kennedy (1961). Calling and associated behavior are treated by Blair (1958, 1959), Gerhardt (1975, 1981), Mecham (1960), Neill (1958), and Oldham and Gerhardt (1975, who comment critically on earlier interpretations of call structure). Recordings of the call are on *Voices of the Night* (Cornell University) and *Sounds of North American Frogs* (Folkways Records FX 6166, New York). Various aspects of life history are discussed by Blem and Miller (1980), Bowler (1977), Caldwell et al. (1980), Lee (1968a), Means and Longdon (1976), Wright (1932), and Wright and Wright (1949). Boyd and Goodyear (1971), Farrell and MacMahon (1969), Hutchison et al. (1968), Jaeger and Hailman (1973), and Sustare (1977) treat physiological topics. Jameson and Richmond (1971), Elias and Shapiro (1957), Gaudin (1974), and Parsons and Williams (1962) deal with morphology.

• ETYMOLOGY. The name *gratiosa* is derived from the Latin *gratiosus*, meaning favored or beloved.





FIGURE. Audiospectrogram of call of *Hyla gratiosa*: 0.5 mi N Hawthorne, Alachua Co., Florida, 23 June 1977, recorded by J. Caldwell; 45 Hz filter left, 300 Hz right.



MAP. Solid circle marks restricted type-locality; open circles indicate other localities; localities outside of shaded area represent disjunct populations.

LITERATURE CITED

- Altig, Ronald. 1970. A key to the tadpoles of the continental United States and Canada. *Herpetologica* 26(2):180-207.
- 1972. Notes on the larvae and premetamorphic tadpoles of four *Hyla* and three *Rana* with notes on tadpole color patterns. *J. Elisha Mitchell Sci. Soc.* 88:113-119.
- , and William L. Pace. 1974. Scanning electron photomicrographs of tadpole labial teeth. *J. Herpetol.* 8(3):247-251.
- Behler, John L., and F. Wayne King. 1979. The Audubon Society field guide to North American reptiles and amphibians. Alfred A. Knopf, New York. 719 p.
- Black, Irving H., and Kenneth L. Gosner. 1958. The barking tree frog, *Hyla gratiosa*, in New Jersey. *Herpetologica* 13(4):254-255.
- Blair, W. Frank. 1958. Call difference as an isolation mechanism in Florida species of hylid frogs. *Quart. J. Florida Acad. Sci.* 21(1):32-48.
- 1959. Call structure and species groups in U.S. treefrogs (*Hyla*). *Southwestern Natur.* 3(1-4):77-89.
- Blair, W. F., A. P. Blair, P. Brodtkorb, F. R. Cagle, and G. A. Moore. 1957. *Vertebrates of the United States*. McGraw-Hill, New York. 819 p.
- Blaney, Richard M. 1971. An annotated check list and biogeographic analysis of the insular herpetofauna of the Apalachicola region, Florida. *Herpetologica* 27(4):406-430.
- Blem, Charles R., and M. A. Miller. 1980. The barking treefrog. *Virginia Wildlife* 41:16-17.
- Bowler, J. Kevin. 1977. Longevity of reptiles and amphibians in North American collections as of 1 November, 1975. *Soc. Stud. Amphib. Rept. Misc. Publ., Herpetol. Circ.* (6):iv + 32.
- Boyd, Claude E., and C. Philip Goodyear. 1971. The protein content of some common reptiles and amphibians. *Herpetologica* 27(3):317-320.
- Caldwell, J. P., J. H. Thorp, and T. O. Jervey. 1980. Predator-prey relationships among larval dragonflies, salamanders, and frogs. *Oecologia* 46:285-289.
- Carr, Archie F., and Coleman J. Goin. 1955. *Guide to the reptiles, amphibians and fresh-water fishes of Florida*. Univ. Florida Press, Gainesville. ix + 341 p.
- Chantell, Charles J. 1964. Some Mio-Pliocene hylids from the Valentine formation of Nebraska. *Amer. Midland Natur.* 72(1):211-225.
- Cochran, Doris M. 1961. Type specimens of reptiles and amphibians in the United States National Museum. *U.S. Nat. Mus. Bull.* (220):xv + 1-289.
- Conant, Roger. 1975. *A field guide to reptiles and amphibians of eastern and central North America*. Second edition. Houghton Mifflin Co., Boston. xviii + 429 p.
- Cope, Edward D. 1885. A contribution to the herpetology of Mexico. I. The collection of the Comision Cientifica. *Proc. Amer. Phil. Soc.* 22(4):379-404.
- 1889. The Batrachia of North America. *U.S. Nat. Mus. Bull.* (34):3-525.
- Elias, Hans, and Jack Shapiro. 1957. Histology of the skin of some toads and frogs. *Amer. Mus. Novitates* (1819):1-27.
- Farrell, Michael P., and James A. MacMahon. 1969. An eco-physiological study of water economy in eight species of tree frogs (Hylidae). *Herpetologica* 25(4):279-294.
- Fortman, Jon, and Ronald Altig. 1973. Characters of F₁ hybrid tadpoles between six species of *Hyla*. *Copeia* 1973(3):411-416.
- , and — 1974. Characters of F₁ hybrid frogs from six species of *Hyla* (Anura:Hylidae). *Herpetologica* 30(3):221-234.
- Gaudin, Anthony J. 1974. An osteological analysis of Holarctic tree frogs, family Hylidae. *J. Herpetol.* 8(2):141-152.
- Gerhardt, H. Carl. 1974. The vocalizations of some hybrid treefrogs: acoustic and behavioral analyses. *Behaviour* 49(1-2):130-151.
- 1975. Sound pressure levels and radiation patterns of the vocalizations of some North American frogs and toads. *J. Comp. Physiol.* 102(1):1-12.
- 1981. Mating call recognition in the barking treefrog (*Hyla gratiosa*): responses to synthetic calls and comparisons with the green treefrog (*Hyla cinerea*). *Ibid.* 144:17-25.
- , Sheldon I. Guttman, and Alvan A. Karlin. 1980. Natural hybrids between *Hyla cinerea* and *Hyla gratiosa*: morphology, vocalization and electrophoretic analysis. *Copeia* 1980(4):577-584.
- Gibbons, J. Whitfield, and J. W. Coker. 1978. Herpetofaunal colonization patterns of Atlantic Coast barrier islands. *Amer. Midland Natur.* 99(1):219-233.
- Hilton, William A. 1954. The chorioid plexuses of frogs and toads. *Herpetologica* 10(2):111-114.
- Hutchison, Victor H., Walter E. Whitford, and Margaret Kohl. 1968. Relation of body size and surface area to gas exchange in anurans. *Physiol. Zool.* 41(1):65-85.
- Jaeger, Robert G., and Jack P. Hailman. 1973. Effects of intensity on the phototactic responses of adult anuran amphibians: A comparative study. *Z. Tierpsychol.* 33:352-407.
- Jacob, James S. 1980. Geographic distribution: *Hyla gratiosa*. *SSAR Herpetol. Rev.* 11(2):36.
- Jameson, David L., and R. C. Richmond. 1971. Parallelism and convergence in the evolution of size and shape in Holarctic *Hyla*. *Evolution* 25(3):497-508.
- LeConte, John. 1856. Description of a new species of *Hyla* from Georgia. *Proc. Acad. Natur. Sci. Philadelphia* 8:146.
- Lee, David S. 1968a. Herpetofauna associated with central Florida mammals. *Herpetologica* 24(1):83-84.
- 1968b. Observations on hybrid *Hyla gratiosa* × *cinerea* in central Florida. *Bull. Maryland Herpetol. Soc.* 4(4):76-78.
- Livezey, Robert L., and Albert H. Wright. 1947. A synoptic key to the salientian eggs of the United States. *Amer. Midland Natur.* 37(1):179-227.
- Malnate, Edmond V. 1971. A catalog of primary types in the herpetological collections of the Academy of Natural Sciences, Philadelphia (ANSP). *Proc. Acad. Natur. Sci. Philadelphia* 123(9):345-375.
- Martof, Bernard S. 1963. Some observations on the herpetofauna of Sapelo Island, Georgia. *Herpetologica* 19(1):70-72.
- Means, D. Bruce, and Clive J. Longden. 1976. Aspects of the biology and zoogeography of the pine barrens treefrog (*Hyla andersoni*) in northern Florida. *Herpetologica* 32(2):117-130.
- Mecham, John S. 1960. Introgressive hybridization between two southeastern treefrogs. *Evolution* 14(4):445-451.
- 1965. Genetic relationships and reproductive isolation in southeastern frogs of the genera *Pseudacris* and *Hyla*. *Amer. Midland Natur.* 74(2):269-308.
- Monroe, Burt L., Jr., and Raymond W. Giannini. 1977. Distribution of the barking treefrog in Kentucky. *Trans. Kentucky Acad. Sci.* 38(3-4):143-144.
- , and Ralph W. Taylor. 1972. Occurrence of the barking treefrog, *Hyla gratiosa*, in Kentucky. *J. Herpetol.* 6(1):78.
- Moore, John A. 1955. Abnormal combinations of nuclear and cytoplasmic systems in frogs and toads. *Advan. Genet.* 7:139-182.
- Mount, Robert H. 1975. *The reptiles and amphibians of Alabama*. Auburn Univ. Agr. Exp. Sta., Auburn, Alabama. vii + 347 p.
- Neill, Wilfred T. 1958. The varied calls of the barking treefrog, *Hyla gratiosa* LeConte. *Copeia* 1958(1):44-46.
- Oldham, Robert S., and H. Carl Gerhardt. 1975. Behavioral isolating mechanisms of the treefrogs *Hyla cinerea* and *H. gratiosa*. *Copeia* 1975(2):223-231.
- Parsons, Thomas, and Ernest E. Williams. 1962. The teeth of Amphibia and their relation to amphibian phylogeny. *J. Morph.* 110(3):375-390.
- Pyburn, William, and J. P. Kennedy. 1961. Hybridization in U.S. treefrogs of the genus *Hyla*. *Proc. Biol. Soc. Washington* 74:157-160.
- Rageot, Roger H. de, A. Damalas, T. Gibbon, and B. Fox. 1969. Observations regarding three rare amphibians in Surry Co. Virginia. *Herpetol. Soc. Bull.* (63):3-5.
- Rossman, Douglas A. 1958. The treefrog *Hyla gratiosa* in Tennessee. *Herpetologica* 14(1):40.
- Schmidt, Karl P. 1953. A check list of North American amphibians and reptiles. Sixth ed. *Amer. Soc. Ichthyol. Herpetol.* viii + 280 p.
- Sustare, B. D. 1977. Characterizing parameters of response to light intensity for six species of frogs. *Behav. Processes* 2(1977):101-112.
- Wright, Albert H. 1929. Synopsis and description of North American tadpoles. *Proc. United States Nat. Mus.* 74(11):1-70.
- 1932. Life-histories of the frogs of Okefinokee Swamp, Georgia. *North American Salientia (Anura)* No. 2. Macmillan, New York. xv + 497 p.
- , and A. A. Wright. 1949. *Handbook of frogs and toads of the United States and Canada*. Third edition. Comstock Publ. Co., Ithaca, New York. xii + 640 p.

JANALEE P. CALDWELL, EMORY UNIVERSITY, ATLANTA, GEORGIA 30322.

Primary editor for this account, Richard G. Zweifel.

Published 16 September 1982 and Copyright 1982 by the SOCIETY FOR THE STUDY OF AMPHIBIANS AND REPTILES.